-continued

85		90	95
Ala Arg Asp Pro Phe 100	Leu His Phe Trp 105		eu Val Thi .0
<pre><210> SEQ ID NO 23 <211> LENGTH: 15 <212> TYPE: PRT <213> ORGANISM: Art <220> FEATURE: <223> OTHER INFORMA</pre>	-	3	
<400> SEQUENCE: 23			
Gly Gly Gly Gly Ser 1 5	Gly Gly Gly Gly	Ser Gly Gly Gly G	y Ser 15

What is claimed is:

- 1. An antibody or an antigen-binding fragment thereof that binds to Ep-CAM, wherein the heavy chain variable CDR3 region has the amino acid sequence SEQ ID NO: 6, 7, 8, 3, 5, or 9.
- 2. The antibody or the antigen-binding fragment thereof of claim 1, wherein the heavy chain variable CDR1 has the amino acid sequence of SEQ ID NO: 10, and the heavy chain variable CDR2 has the amino acid sequence of SEQ ID NO: 11.
- 3. The antibody or the antigen-binding fragment thereof of claim 1, wherein the light chain variable CDR1 has the amino acid sequence of SEQ ID NO: 12, the light chain variable CDR2 has the amino acid sequence of SEQ ID NO: 13, and the light chain variable CDR3 has the amino acid sequence of SEQ ID NO: 14.
- **4**. The antibody or the antigen-binding fragment thereof of claim **1**, wherein V_L comprises the amino acid sequence of SEQ ID NO: 15 or 16, and V_H comprises the amino acid sequence of SEQ ID NO: 17, 18, 19, 20, 21, or 22.
- **5**. The antigen-binding fragment of claim **1**, which is single chain variable fragment (scFv).
- **6**. The scFv of claim **5**, wherein the heavy chain variable CDR3 region has the amino acid sequence SEQ ID NO: 6, 7, or 8.
- 7. A chimeric antigen receptor fusion protein (CAR) comprising from N-terminus to C-terminus:

- (i) the scFv of claim 6,
- (ii) a hinge domain
- (iii) a transmembrane domain,
- (iv) at least one co-stimulatory domains, and
- (v) an activating domain.
- **8**. The CAR according to claim **7**, wherein the costimulatory domain is selected from the group consisting of CD28, 4-1BB, ICOS-1, CD27, OX-40, GITR, and DAP10.
- **9**. The CAR according to claim **7**, wherein the activating domain is CD**3** zeta.
- 10. An isolated nucleic acid sequence encoding the CAR of claim 7.
- 11. T cells or natural killer cells modified to express the CAR of claim 7.
- 12. An adoptive cell therapy method for treating cancer, comprising the steps of:
 - administering the CAR-T cells of claim 11 to a subject suffering from cancer, wherein the cancer cells of the subject overexpress EpCAM, and the CAR T cells bind to the cancer cells to kill the cancer cells.
- 13. The method according to claim 12, wherein the cancer is colon, intestine, breast, lung, prostate, gastric, pancreas, bladder, gall bladder, nasopharyngeal, colorectal, ovarian, or lung cancer.

* * * * *